

See discussions, stats, and author profiles for this publication at: <http://www.researchgate.net/publication/261476577>

# Stoichiometric regulation of phytoplankton toxins

ARTICLE *in* ECOLOGY LETTERS · APRIL 2014

Impact Factor: 13.04 · DOI: 10.1111/ele.12280

---

CITATIONS

5

---

DOWNLOADS

104

---

VIEWS

93

5 AUTHORS, INCLUDING:



**Dedmer Van de Waal**

Netherlands Institute of Ecology (NIOO-KN...

20 PUBLICATIONS 203 CITATIONS

SEE PROFILE



**Val H. Smith**

University of Kansas

109 PUBLICATIONS 8,619 CITATIONS

SEE PROFILE

## LETTER

## Stoichiometric regulation of phytoplankton toxins

Dedmer B. Van de Waal,<sup>1\*</sup>  
Val H. Smith,<sup>2</sup> Steven A. J.  
Declerck,<sup>1</sup> Eva C. M. Stam<sup>1</sup> and  
James J. Elser<sup>3</sup>

### Abstract

Ecological Stoichiometry theory predicts that the production, elemental structure and cellular content of biomolecules should depend on the relative availability of resources and the elemental composition of their producer organism. We review the extent to which carbon- and nitrogen-rich phytoplankton toxins are regulated by nutrient limitation and cellular stoichiometry. Consistent with theory, we show that nitrogen limitation causes a reduction in the cellular quota of nitrogen-rich toxins, while phosphorus limitation causes an increase in the most nitrogen-rich paralytic shellfish poisoning toxin. In addition, we show that the cellular content of nitrogen-rich toxins increases with increasing cellular N : P ratios. Also consistent with theory, limitation by either nitrogen or phosphorus promotes the C-rich toxin cell quota or toxicity of phytoplankton cells. These observed relationships may assist in predicting and managing toxin-producing phytoplankton blooms. Such a stoichiometric regulation of toxins is likely not restricted to phytoplankton, and may well apply to carbon- and nitrogen-rich secondary metabolites produced by bacteria, fungi and plants.

### Keywords

Ecological stoichiometry, harmful algal blooms, harmful cyanobacteria, natural toxins, nutrient limitation, secondary metabolites.

*Ecology Letters* (2014)

### INTRODUCTION

It is well established that the elemental composition of both prokaryotic and eukaryotic cells strongly depends on the supplies of chemical elements, such as the key nutrients nitrogen (N) and phosphorus (P) (Sterner & Elser 2002; Klausmeier *et al.* 2004). Primary producers use light energy to convert CO<sub>2</sub> and nutrients into an enormous variety of organic compounds, including fatty acids, amino acids and nucleic acids. These three key groups of biomolecules differ significantly in their elemental composition, notably in their relative carbon (C), N and P contents. Similarly, there are major differences in the C : N : P stoichiometry of secondary cellular metabolites, which are known to serve as signalling molecules as well as deterrents against pathogens and herbivores (Keller & Surette 2006; Iason *et al.* 2013). Some of the secondary metabolites produced by planktonic aquatic primary producers (unicellular and colonial cyanobacteria and eukaryotic algae, collectively termed 'phytoplankton' hereafter) are toxic to invertebrates, fish, birds and mammals. Blooms of such toxic phytoplankton are of increasing global concern because their appearance and proliferation are driven by high anthropogenic loadings of N and P to surface waters (eutrophication) (Anderson *et al.* 2002; Smith 2003). Thus, there is a strong need to gain a better understanding of how the absolute and relative supplies of key nutrients affect the production of toxic compounds by bloom-forming phytoplankton.

Toxins produced by phytoplankton are stoichiometrically diverse, ranging from N-rich alkaloids to C-based polyketides (Table 1). Previous studies involving particular phytoplankton taxa have demonstrated that the production and cellular contents (internal cell 'quota') of C- and N-rich toxins respond strongly to the relative availability of nutrients (Granéli *et al.* 1998; Sivonen & Jones 1999; Granéli & Flynn 2006; Van de Waal *et al.* 2009). Here, we present the first quantitative analysis of the regulation of phytoplankton toxins by nutrient limitation. We analysed >500 published studies and compiled a total of 180 datasets, covering 31 phytoplankton species representing 19 different genera (Fig. S1; Table S1). Using these data, we analysed the toxin cell quotas or toxicities of phytoplankton cells grown under either N- or P-limitation, and we compared these responses with cells grown in nutrient-sufficient controls. Based on stoichiometric theory (Sterner & Elser 2002) and current understanding of phytoplankton physiology, we predicted that production of C-rich toxic compounds would be elevated under both N- and P-limitation because, under such conditions, cellular C is by definition in excess and can accumulate in secondary pools. Following similar reasoning, we also predicted that production of N-rich toxins would be inhibited under N-limitation (because N is shunted into intracellular pools to support active growth), but amplified under P-limitation (because N is in excess and accumulates in secondary intracellular pools).

<sup>1</sup>Department of Aquatic Ecology, Netherlands Institute of Ecology (NIOO-KNAW), Post Office Box 50, Wageningen, 6700 AB, The Netherlands

<sup>2</sup>Department of Ecology and Evolutionary Biology, University of Kansas, Lawrence, Kansas, 66045, USA

<sup>3</sup>School of Life Sciences, Arizona State University, Tempe, AZ, 85287-4501, USA

\*Correspondence: E-mail: d.vandewaal@nioo.knaw.nl

**Table 1** Overview of toxins grouped according to their C : N ratio.

Group	Toxins	Molecular formula	C : N ratio*	
N-rich	PSP	Paralytic Shellfish Poisoning toxins (incl. saxitoxin, STX)	C <sub>10</sub> H <sub>19</sub> N <sub>7</sub> O <sub>4</sub>	1.5
	CYN	Cylindrospermopsin	C <sub>15</sub> H <sub>21</sub> N <sub>5</sub> O <sub>7</sub> S	3.0
	MC	Microcystin (incl. microcystin-LR; microcystin-RR)	C <sub>49</sub> H <sub>74</sub> N <sub>10</sub> O <sub>12</sub> (MC-LR); C <sub>49</sub> H <sub>75</sub> N <sub>13</sub> O <sub>12</sub> (MC-RR)	4.3
	NOD	Nodularin	C <sub>41</sub> H <sub>60</sub> N <sub>8</sub> O <sub>10</sub>	5.1
C-rich	ANTX	Anatoxin a	C <sub>10</sub> H <sub>15</sub> NO	10
	ASP	Amnesic Shellfish Poisoning toxin (incl. domoic acid, DA)	C <sub>15</sub> H <sub>21</sub> NO <sub>6</sub>	15
	GYM	Gymnodimine	C <sub>32</sub> H <sub>45</sub> NO <sub>4</sub>	32
	SPX	Spirolide	C <sub>42</sub> H <sub>61</sub> NO <sub>7</sub>	42
	PLTX	Palytoxin	C <sub>129</sub> H <sub>223</sub> N <sub>3</sub> O <sub>54</sub>	43
	OVTX	Ovatoxin (incl. OVTX-a)	C <sub>129</sub> H <sub>223</sub> N <sub>3</sub> O <sub>52</sub>	43
	DSP	Diarrhoeic Shellfish Poisoning toxins (incl. okadaic acid, OA; dinophysins toxins, DTX-1, DTX-2)	C <sub>44</sub> H <sub>68</sub> O <sub>13</sub> (OA, DTX-2); C <sub>45</sub> H <sub>70</sub> O <sub>13</sub> (DTX-1);	-
	NSP	Neurotoxic Shellfish Poisoning toxin (incl. brevetoxin, PbTx)	C <sub>49</sub> H <sub>70</sub> O <sub>13</sub>	-
	KMTX	Karlotoxins (incl. KMTX-1-1, KMTX-2)	C <sub>67</sub> H <sub>120</sub> O <sub>24</sub> (KMTX-1-1); C <sub>67</sub> H <sub>121</sub> ClO <sub>24</sub> (KMTX-2)	-
	CTX	Ciguatoxin	C <sub>61</sub> H <sub>88</sub> O <sub>19</sub>	-
	MTX	Maitotoxin	C <sub>164</sub> H <sub>256</sub> O <sub>68</sub> S <sub>2</sub> Na <sub>2</sub>	-
	HA	Haemolytic activity based on bioassays		-

\*Values refer to the described toxins, or to the average of the shown analogues.

## MATERIAL AND METHODS

Data were obtained by ISI Web of Science searches with the terms: ('phytoplankton' or 'cyanobacteri\*' or 'dinoflagellate\*') and '\*toxin\*' and ('nutrient\*' or 'nitr\*' or 'phosph\*') (Fig. S1; Tables S1 and S2). Estimates of toxin quota or toxicity of phytoplankton cells grown under low N and/or low P were recovered from the stationary growth phase of batch culture experiments, and from steady-state conditions of continuous cultures. Toxin quotas or toxicities of cells grown under nutrient-sufficient conditions, or at maximum growth rates during the exponential phase, served as controls. Studies reporting earlier published data were not included. We made a distinction between N-rich and C-rich toxins based on elemental composition. Toxins with a molar C : N ratio <6.6 (i.e. less than the Redfield ratio) were considered 'N rich', whereas toxins with a higher molar C : N ratio were considered 'C rich' (Table 1). For each study, we took toxin quotas or toxicities and calculated the average response ratio (RR) by normalising toxin quotas or toxicities from the low N and low P conditions to the control. Furthermore, we derived cellular N : P ratios expressed as molar ratios.

Toxins were found to be represented by varying numbers of studies (Fig. S1). Some toxins were produced by multiple species, and these toxins were represented by different numbers of species (Fig. S1). To correct for biases that may result from such unbalanced and nested data structure, we worked with weighted means and variances. First, we calculated the average response and variance for each toxin that was represented by at least three or more independent studies (i.e.  $n > 3$ ). In case, a toxin was produced by multiple species, and one species was represented by more than two studies, we first calculated the mean of this species and then calculated the average response of the toxin across species. In case, multiple species were represented by multiple studies, the weighted average response of a toxin across species could be calculated according to (Borenstein *et al.* 2009):

$$M_{RR,W} = \frac{\sum_{i=1}^n M_{RRi} W_i}{\sum_{i=1}^n W_i} \quad (1)$$

with  $M_{RRi}$  being the mean response ratio for species  $i$  and weight  $W_i$  being the reciprocal of the variance  $V_i$ , i.e. the variance among studies for each species  $i$ , calculated as:

$$V_i = \frac{1}{m-1} \sum_{j=1}^m (RR_{ij} - M_{RRi})^2 \quad (2)$$

Similarly, the overall weighted average response of toxins across the means of all N- or C-rich toxins was calculated. We assessed the statistical significance of differences between the calculated RRs and their respective controls with a one sample  $t$ -test comparing the weighted mean RR with 1, which represents the absence of a response ( $\alpha < 0.05$ ).


## RESULTS

The assembled studies yielded 100 data sets with N-rich toxins (49 with N-limitation and 51 with P-limitation) and 80 data sets with C-rich toxins (37 with N-limitation and 43 with P-limitation; Fig. S1). N-rich toxins predominantly consisted of Paralytic Shellfish Poisoning (PSP) toxins produced by marine dinoflagellates ( $n = 57$ ), followed by microcystins (MCs) produced by freshwater cyanobacteria ( $n = 34$ ). Data on C-rich toxins comprised 12 compounds, groups of compounds, or toxicities associated with C-rich toxins. The C-rich toxins were almost exclusively found in marine (including brackish water) ecosystems and consisted of various toxins produced by dinoflagellates (including Diarrhoeic Shellfish Poisoning (DSP) toxins,  $n = 18$ , karlotoxins (KMTX),  $n = 11$  and Neurotoxic Shellfish Poisoning (NSP) toxins,  $n = 10$ ), diatoms (Amnesic Shellfish Poisoning (ASP) toxins,  $n = 9$ ), and prymnesiophytes (haemolytic activity (HA),  $n = 12$ ). Various toxins were only represented by a limited number of studies ( $n < 3$ ) and were therefore excluded from our analyses. For

most of the toxins, experiments with N-limitation and P-limitation tended to be equally well represented (Fig. S1).

The impact of nutrient limitation on toxin quota or toxicity varied greatly and depended strongly on the stoichiometric nature of the toxins. Consistent with theoretical predictions, our analysis revealed that N-rich toxins generally decreased upon N-limitation while C-rich toxins generally increased upon limitation by either N or P (Figs 1 and 2). More specifically, cellular quota of both N-rich PSP toxins and MCs decreased significantly upon N-limitation, irrespective of the producing taxonomic group. Upon P-limitation, cellular quota of dinoflagellate PSP toxins increased by 32%, while MCs decreased by 75% (Fig. 1; Table 2). Relative changes in C-rich toxins were generally larger than the N-rich toxins, particularly under P-limitation, although variances within toxins were also higher. Significant increases in cellular toxin quota ranged from 99% in NSP toxins under N-limitation to 215% in KMTX under P-limitation (Fig. 2; Table 2).

Under balanced, non-limiting growth conditions, molar phytoplankton N : P ratios typically converge upon the so-called 'Redfield ratio' of N : P = 16 (Sterner & Elser 2002).



**Figure 1** Relative changes of N-rich toxins in response to nutrient limitation. (a) N-limitation and (b) P-limitation. White symbols show weighted means for individual toxins with error bars indicating SE across means of species; grey symbols show the overall weighted means of N-rich toxins with error bars indicating SE across means of toxins. The horizontal line indicates absence of response. Asterisks indicate significances  $P < 0.01$  (\*\*) and  $P < 0.05$  (\*).

**Table 2** Results of one-sample t-tests comparing relative changes in N- and C-rich toxin quotas or toxicities between a treatment and its control, with sample size ( $n$ ), mean change, 95% confidence interval and  $P$ -value at a significance level of 0.05 (indicated in bold).

	$n$	Mean	Variance	95% CI	$P$ -value
N-rich					
N-limitation					
PSP	5*	0.65	0.009	0.46–0.83	<b>0.011</b>
MC	2*	0.49	0.003	0.38–0.60	<b>0.035</b>
All	2†	0.53	0.002	0.44–0.62	<b>0.033</b>
P-limitation					
PSP	5*	1.32	0.004	1.19–1.44	<b>0.004</b>
MC	3*	0.25	0.008	0.08–0.43	<b>0.008</b>
NOD	5	0.93	0.886	0.11–1.76	0.440
All	3†	0.96	0.003	0.86–1.06	0.271
C-rich					
N-limitation					
DSP	2*	1.69	1.95	–1.04–4.43	0.353
KMTX	5	2.77	0.93	1.92–3.61	<b>0.008</b>
NSP	5	1.99	0.15	1.65–2.33	<b>0.002</b>
ASP	2*	1.04	1.02	–0.37–2.44	0.484
HA	2*	4.41	6.47	0.88–7.93	0.155
All	5†	2.00	0.11	1.36–2.64	<b>0.019</b>
P-limitation					
DSP	2*	2.04	4.40	–0.34–4.41	0.241
KMTX	6	3.15	3.72	1.61–4.70	<b>0.021</b>
NSP	5	2.30	0.34	1.79–2.82	<b>0.004</b>
ASP	2*	6.25	4.87	1.93–10.58	0.127
HA	2*	6.48	25.69	0.74–12.21	0.101
ANTX	3	1.10	0.16	0.65–1.56	0.351
All	6†	1.66	0.10	1.04–2.29	<b>0.002</b>


\*Analysis based on means or weighted means across species.

†Analysis based on weighted means across toxins.

Indeed, in the control treatments of the experiments we assembled, cellular N : P ratios resembled the Redfield ratio, whereas cellular N : P stoichiometry was almost always lower than Redfield under N-limitation and higher than Redfield under P-limitation (Fig. 3). N-rich toxins showed a positive linear relationship with cellular N : P ratios. The response ratios of the N-rich PSP toxins and MCs decreased below 1 at N : P ratios below the Redfield ratio, but increased beyond 1 when N : P ratios exceeded the Redfield ratio (Fig. 3a). In contrast, the C-rich DSP toxins and HA increased at N : P ratios both below and above the Redfield ratio (Fig. 3b).

## DISCUSSION


Our results demonstrate that the synthesis of N-rich and C-rich toxins is regulated by the relative availabilities of N and P (Figs. 1 and 2). Upon N-limitation, cellular quota of the N-rich PSP toxins and microcystins decreased, while upon P-limitation, cellular quota of the most N-rich PSP toxins increased. Cellular quota of microcystin, however, showed a strong decrease, and nodularin did not change upon P-limitation (Fig. 1). These contrasting responses to P-limitation may reflect the fact that, unlike eukaryotic algae, cyanobacteria can shunt excess N into the storage compound cyanophycin (Allen 1984), a process that likely prevents N from accumulat-



**Figure 2** Relative changes of C-rich toxins in response to nutrient limitation. (a) N-limitation and (b) P-limitation. White symbols show weighted means for individual toxins with error bars indicating SE across means of species; grey symbols show the overall weighted means of C-rich toxins with error bars indicating SE across means of toxins. The horizontal line indicates absence of response. Asterisks indicate significances  $P < 0.01$  (\*\*) and  $P < 0.05$  (\*).

ing primarily as microcystin. C-rich toxins generally increased upon limitation by both N and P (Fig. 2). Most of the C-rich toxins were produced by eukaryotic algae and are polyketides, whose synthesis is tightly coupled with cellular C acquisition (Staunton & Weissman 2001). Thus, under conditions of nutrient limitation, the relative excess of energy and newly synthesised organic C cannot be used for cell growth and instead appears to be shunted into C-rich molecules such as lipids and polyketides, including those that have toxic effects. We also observed a general increase in N-rich toxins with cellular N : P ratio (Fig. 3a), whereas C-rich toxins showed an initial decrease with N : P ratios until reaching the Redfield ratio, after which their content increased (Fig. 3b). These findings are consistent with the 'V-shaped' response to N : P ratios that has been reported previously for haemolytic activity associated with C-rich compounds (Granéli & Flynn 2006; Granéli *et al.* 2012).

While most of the selected data sets reported cellular toxin quotas, a few (~6%) assessed toxicities relative to a standard toxin analogue, for instance, as saxitoxin equivalents (Table S1). Changes in these relative toxicities are generally associated with changes in cellular toxin quota, yet may also relate to a shift in the toxin composition (Cembella 1998; Sivonen & Jones



**Figure 3** Relative changes of N- and C-rich toxins in response to cellular N:P ratios. (a) N-rich toxins and (b) C-rich toxins. Symbols show weighted means for individual toxins with error bars indicating SE across means of species. The horizontal line indicates an absence of response, and the vertical line illustrates the Redfield N : P ratio (N : P = 16). The solid grey line in (a) shows the linear fit to all data ( $R^2 = 0.989$ ,  $n = 6$ ,  $P < 0.001$ ).

1999). Because the few selected studies reporting relative toxicities generally showed none, or only minor shifts in toxin composition, we considered the observed changes in relative toxicities to mainly reflect changes in the N-rich toxin quota. All reported haemolytic activities of prymnesiophytes were assessed by bioassays and have been ascribed to various C-rich compounds, such as glycolipids and the polyether prymnesin (Edvardsen & Imai 2006; Granéli *et al.* 2012). Although haemolytic activity showed a high average response to nutrient limitation, the variance was also very high (Table 2), which may reflect a higher inherent variability of bioassay data relative to direct chemical measurements of the cellular quota of a specific compound. Nevertheless, our analysis reveals that the response of haemolytic activity by prymnesiophytes towards both N- and P-limitation is consistent with the general response of C-rich toxins, and therefore presumably reflects changes in cellular quotas of a C-rich compound.

Toxins produced by various harmful phytoplankton taxonomic groups have been shown to exhibit a broad range of potential functions, including grazer deterrence, prey capture and resource uptake (see also Granéli & Turner 2006; Glibert

& Burkholder 2011). The question remains whether our results reflect a possible role of toxins in acquiring N and P, or if the toxin quota merely responds to the relative availability of N and C within the cells. Production of N-rich toxins generally decreased with N-limitation, suggesting a cost rather than a benefit under N-limitation. Cellular contents of N-rich cyanobacterial toxins strongly decreased or remained unaltered upon P-limitation, whereas the quota of N-rich dinoflagellate PSP toxins increased. C-rich toxins in dinoflagellates and diatoms, as well as haemolytic activity caused by prymnesiophytes, also increased with P-limitation, and the dinoflagellate karlotoxin and NSP toxins increased significantly with N-limitation. These findings suggest that toxin synthesis may be related to resource acquisition, specifically C-rich toxins for acquiring P. Indeed, C-rich toxins produced by prymnesiophytes seem to be involved in prey capture (Skovgaard *et al.* 2003; Tillmann 2003). Furthermore, nutrient limitation may stimulate feeding by mixotrophic dinoflagellates and prymnesiophytes (Legrand *et al.* 2001; Smalley *et al.* 2003), although it is not clear whether this is related to their toxicity (Stoecker *et al.* 2006). Thus, whether toxins stimulated by nutrient limitation are indeed involved in resource acquisition and prey capture, or whether they merely respond to a relative excess of non-limiting resources required for their synthesis, needs to be further elucidated.

Our findings illustrate that the extent to which phytoplankton are able to produce toxins, and thereby exploit their putative function, will partially depend on the relative availabilities of N and P in the environment. Because N and P often limit phytoplankton growth in both freshwater and marine ecosystems (Elser *et al.* 2007), the cellular production of C-rich toxins avoids resource allocation conflicts between population growth and secondary metabolite synthesis. Production of N-rich toxins, however, does decrease with N-limitation, suggesting that N is allocated to growth rather than secondary metabolites. Growth and the synthesis of biochemicals together will determine the elemental stoichiometry of phytoplankton, which shows distinct differences between taxonomic groups. For instance, dinoflagellates exhibit higher C : N and C : P ratios as compared to prymnesiophytes and diatoms (Quigg *et al.* 2003), whereas cyanobacteria have generally higher optimal N : P ratios than diatoms (Hillebrand *et al.* 2013). Most of the C-rich toxins studied here were indeed produced by dinoflagellates (i.e. 10 of 12 toxins; Fig. S1), and the reported N-rich toxins were produced by cyanobacteria (i.e. MC, PSP, NOD and CYN) or seem to have a common origin with cyanobacteria (i.e. PSP; Moustafa *et al.* 2009; Hackett *et al.* 2013). We did not observe differences in the elemental composition between taxonomic groups, yet our results demonstrate that the cellular quota of the N-rich PSP toxins and microcystins are strongly correlated with N : P stoichiometry, and this relationship is irrespective of the producing taxonomic group, both at the genus as well as at the species level (Fig. 3). To what extent the stoichiometry of a taxonomic group may influence the stoichiometry of the toxins that can be produced remains to be elucidated. Obviously, cellular elemental ratios strongly reflect the intracellular content of major biochemicals such as proteins and nucleic acids (Sterner & Elser 2002). Here, we show that they

may also determine, at least partially, the stoichiometry of secondary metabolites that can be synthesised by the cell.

The strong association of phytoplankton toxins with N : P stoichiometry has important implications for interpreting and predicting the responses of phytoplankton-associated toxins to human-induced changes in the magnitude and ratios of N and P supplies to surface waters. One of the clearest and most predictable responses of both freshwater and marine ecosystems to increased nutrient loading is the profound stimulation of phytoplankton biomass, often by orders of magnitude (Smith 2003; Dolman *et al.* 2012; Orihel *et al.* 2012). Total toxin concentrations in natural waters will therefore tend to increase with eutrophication-driven increases in total phytoplankton biomass, but the magnitude of this response will be amplified if nutrient enrichment also selects for dominance by toxin-producing taxa (Anderson *et al.* 2002; Heisler *et al.* 2008; Dolman *et al.* 2012; Scott *et al.* 2013) or clones (Kardinaal *et al.* 2007; Touzet *et al.* 2007) and if nutrient limitation promotes the internal accumulation of toxic molecules, as shown in our analysis. These insights also provide a means for anticipating which toxins will be triggered as a function of N- or P-limitation during bloom termination. Thus, the stoichiometric relationships we have uncovered may provide a practical basis for better forecasting the potential occurrence of toxic compounds in the aquatic environment, possibly allowing the creation of an 'early warning system' for more detailed chemical analyses of suspended particulate matter, both in drinking water supplies and in surface waters that support fisheries and heavy recreational use.

Our analysis demonstrates that the production of stoichiometrically distinct toxins can be explained, at least partially, by the availabilities of key limiting nutrients in the environment and within the cell. However, we note that the production of toxins may also be affected by other environmental factors. In particular, C-rich toxins may be sensitive to changes in CO<sub>2</sub> and light availability, as these resources provide phytoplankton with C and energy. Indeed, enhanced CO<sub>2</sub> availabilities have been shown to increase the production of C-rich toxins such as domoic acid (Sun *et al.* 2011) and karlotoxin (Fu *et al.* 2010), particularly under P-limited conditions. However, there seems no general response of C-rich toxins to increasing light intensities. More specifically, increasing light intensities were shown to cause an increase (Bates 1998; Cusack *et al.* 2002) or a decrease (Auro & Cochlan 2013) in the production of domoic acid, while the sensitivity of haemolytic activity (Baker *et al.* 2007; Granéli *et al.* 2012) and DSP toxins (Nielsen *et al.* 2013) to light intensity seems to be marginal or even absent. Enhanced light intensities were shown to cause an increase in the production of the N-rich PSP toxins (Parkhill & Cembella 1999; Etheridge & Roesler 2005) and microcystins (Wiedner *et al.* 2003; Tonk *et al.* 2005) at limiting light levels, but caused a decrease at saturating light levels. Generally, the response of N- and C-rich toxins to light availability seems less pronounced as compared to nutrient limitation, while enhanced CO<sub>2</sub> availabilities may further promote the production of C-rich toxins under nutrient limiting conditions.

Our results provide a broad picture of how the synthesis of toxins in response to nutrient limitation might be modified by internal allocation processes that shunt excess C and nutrients

not to toxins but towards storage compounds, as in the case of N in cyanobacteria. In higher plants, nutrient limitation may result in a surplus of fixed C as well, which can also be allocated either to storage compounds or to C-rich secondary metabolites (Koricheva *et al.* 1998). Likewise, the availability of C and nutrients has been shown to affect synthesis of C- or N-rich secondary metabolites in bacteria and fungi (Sanchez & Demain 2002; Cornforth & Foster 2013). Even though secondary metabolites are highly diverse in function among producers, the synthesis of these molecules generally has a flexible component that at least partially depends on the interplay between their own elemental composition and the relative availability of key resources. Thus, toxin production can be expected to behave in broadly predictable ways across a wide variety of biota.

Ongoing anthropogenic loading of nutrients will lead to deterioration of water quality not only by promoting biomass accumulation of phytoplankton, but by affecting both the quantities and the biochemical composition of the toxins they can produce, especially when nutrient loading leads to strong stoichiometric imbalances of N and P. For instance, nutrient inputs from non-point sources such as atmospheric N deposition or release of sediment-bound P (Carpenter *et al.* 1998; Elser *et al.* 2009) may shift phytoplankton from N- to P-limitation, or vice versa, and thereby alter their toxicity. Thus, application of stoichiometric theory, together with knowledge of the phytoplankton species and the stoichiometric composition of the toxins they produce, may help us predict whether nutrient loading will cause an increase, decrease, or no change in overall toxicity in natural waters (Anderson *et al.* 2002; Heisler *et al.* 2008), with important potential consequences for the various services provided by freshwater and marine ecosystems.

## ACKNOWLEDGEMENTS

We thank the editor and anonymous referees for their valuable comments on the manuscript. This work was supported in part by the NSF grants DMS-0342239 to VHS and DEB-0950179 to JJE.

## STATEMENT OF AUTHORSHIP

D.B.v.d.W. and E.C.M.S. recovered the data. All authors analysed the data and discussed the results. D.B.v.d.W. wrote the first draft of the manuscript, and all authors contributed substantially to revisions.

## REFERENCES

- Allen, M.M. (1984). Cyanobacterial cell inclusions. *Annu. Rev. Microbiol.*, 38, 1–25.
- Anderson, D.M., Glibert, P.M. & Burkholder, J.M. (2002). Harmful algal blooms and eutrophication: nutrient sources, composition, and consequences. *Estuaries*, 25, 704–726.
- Auro, M.E. & Cochlan, W.P. (2013). Nitrogen utilization and toxin production by two diatoms of the *Pseudo-nitzschia pseudodelicatissima* Complex: *P. cuspidata* and *P. fryxelliana*. *J. Phycol.*, 49, 156–169.
- Baker, J.W., Grover, J.P., Brooks, B.W., Urena-Boeck, F., Roelke, D.L., Errera, R. *et al.* (2007). Growth and toxicity of *Prymnesium parvum* (Haptophyta) as a function of salinity, light, and temperature. *J. Phycol.*, 43, 219–227.
- Bates, S.S. (1998). Ecophysiology and Metabolism of ASP Toxin Production. In: *Physiological ecology of harmful algal blooms*. (eds Anderson, D.M., Cembella, A.D., Hallegraeff, G.M.). Springer-Verlag, Berlin Heidelberg, Heidelberg, Germany, pp. 405–426.
- Borenstein, M., Hedges, L.V., Higgins, J.P.T. & Rothstein, H.R. (2009). *Introduction to Meta-Analysis*. John Wiley & Sons, Chichester, UK.
- Carpenter, S.R., Caraco, N.F., Correll, D.L., Howarth, R.W., Sharpley, A.N. & Smith, V.H. (1998). Nonpoint pollution of surface waters with phosphorus and nitrogen. *Ecol. Appl.*, 8, 559–568.
- Cembella, A.D. (1998). Ecophysiology and Metabolism of Paralytic Shellfish Toxins in Marine Microalgae. In: *Physiological Ecology of Harmful Algal Blooms*. (eds Anderson, D.M., Cembella, A.D., Hallegraeff, G.M.). Springer-Verlag, Berlin Heidelberg Heidelberg, Germany, pp. 281–403.
- Cornforth, D.M. & Foster, K.R. (2013). Competition sensing: the social side of bacterial stress responses. *Nat. Rev. Microbiol.*, 11, 285–293.
- Cusack, C.K., Bates, S.S., Quilliam, M.A., Patching, J.W. & Raine, R. (2002). Confirmation of domoic acid production by *Pseudo-nitzschia australis* (Bacillariophyceae) isolated from Irish waters. *J. Phycol.*, 38, 1106–1112.
- Dolman, A.M., Rucker, J., Pick, F.R., Fastner, J., Rohrlack, T., Mischke, U. *et al.* (2012). Cyanobacteria and cyanotoxins: the influence of nitrogen versus phosphorus. *PLoS ONE*, 7, e38757.
- Edwardsen, B. & Imai, I. (2006). The Ecology of Harmful Flagellates within Prymnesiophyceae and Raphidophyceae. In: *Ecology of Harmful Algae*. (eds Granéli, E., Turner, J.T.). Springer-Verlag, Berlin Heidelberg, Heidelberg, Germany, pp. 67–79.
- Elser, J.J., Bracken, M.E.S., Cleland, E.E., Gruner, D.S., Harpole, W.S., Hillebrand, H. *et al.* (2007). Global analysis of nitrogen and phosphorus limitation of primary producers in freshwater, marine and terrestrial ecosystems. *Ecol. Lett.*, 10, 1135–1142.
- Elser, J.J., Andersen, T., Baron, J.S., Bergstrom, A.K., Jansson, M., Kyle, M. *et al.* (2009). Shifts in lake N: P stoichiometry and nutrient limitation driven by atmospheric nitrogen deposition. *Science*, 326, 835–837.
- Etheridge, S.M. & Roesler, C.S. (2005). Effects of temperature, irradiance, and salinity on photosynthesis, growth rates, total toxicity, and toxin composition for *Alexandrium fundyense* isolates from the Gulf of Maine and Bay of Fundy. *Deep Sea Res. (II Top. Stud. Oceanogr.)*, 52, 2491–2500.
- Fu, F.X., Place, A.R., Garcia, N.S. & Hutchins, D.A. (2010). CO<sub>2</sub> and phosphate availability control the toxicity of the harmful bloom dinoflagellate *Karlodinium veneticum*. *Aquat. Microb. Ecol.*, 59, 55–65.
- Glibert, P.M. & Burkholder, J.M. (2011). Harmful algal blooms and eutrophication: 'Strategies' for nutrient uptake and growth outside the Redfield comfort zone. *Chin. J. Oceanol. Limnol.*, 29, 724–738.
- Granéli, E. & Flynn, K. (2006). Chemical and Physical Factors Influencing Toxin Content. In: *Ecology of Harmful Algae*. (eds Granéli, E., Turner, J.T.). Springer-Verlag, Berlin Heidelberg, Heidelberg, Germany, pp. 229–241.
- Granéli, E. & Turner, J.T. (2006). *Ecology of Harmful Algae*. Springer-Verlag, Berlin Heidelberg, Heidelberg, Germany.
- Granéli, E., Johansson, N. & Panosso, R. (1998). Cellular Toxin Contents in Relation to Nutrient Conditions for Different Groups of Phytotoxins. In: *Harmful Algae* (eds Reguera, B., Blanco, J., Fernández, M.L., Wyatt, T.). Xunta de Galicia and IOC-UNESCO, Santiago de Compostella, pp. 321–324.
- Granéli, E., Edwardsen, B., Roelke, D.L. & Hagstrom, J.A. (2012). The ecophysiology and bloom dynamics of *Prymnesium* spp. *Harmful Algae*, 14, 260–270.
- Hackett, J.D., Wisecaver, J.H., Brosnahan, M.L., Kulis, D.M., Anderson, D.M., Bhattacharya, D. *et al.* (2013). Evolution of Saxitoxin Synthesis in Cyanobacteria and Dinoflagellates. *Mol. Biol. Evol.*, 30, 70–78.
- Heisler, J., Glibert, P.M., Burkholder, J.M., Anderson, D.M., Cochlan, W., Dennison, W.C. *et al.* (2008). Eutrophication and harmful algal blooms: a scientific consensus. *Harmful Algae*, 8, 3–13.

- Hillebrand, H., Steinert, G., Boersma, M., Malzahn, A., Meunier, C.L., Plum, C. *et al.* (2013). Golman revisited: faster-growing phytoplankton has lower N:P and lower stoichiometric flexibility. *Limnol. Oceanogr.*, 58, 2076–2088.
- Iason, G.R., Dicke, M. & Hartley, S.E. (2013). *The Ecology of Plant Secondary Metabolites: From Genes to Global Processes*. Cambridge University Press, New York.
- Kardinaal, W.E.A., Janse, I., Kamst-van Agterveld, M., Meima, M., Snoek, J., Mur, L.R. *et al.* (2007). *Microcystis* genotype succession in relation to microcystin concentrations in freshwater lakes. *Aquat. Microb. Ecol.*, 48, 1–12.
- Keller, L. & Surette, M.G. (2006). Communication in bacteria: an ecological and evolutionary perspective. *Nat. Rev. Microbiol.*, 4, 249–258.
- Klausmeier, C.A., Litchman, E., Daufresne, T. & Levin, S.A. (2004). Optimal nitrogen-to-phosphorus stoichiometry of phytoplankton. *Nature*, 429, 171–174.
- Koricheva, J., Larsson, S., Haukioja, E. & Keinanen, M. (1998). Regulation of woody plant secondary metabolism by resource availability: hypothesis testing by means of meta-analysis. *Oikos*, 83, 212–226.
- Legrand, C., Johansson, N., Johnsen, G., Borsheim, K.Y. & Graneli, E. (2001). Phagotrophy and toxicity variation in the mixotrophic *Prymnesium patelliferum* (Haptophyceae). *Limnol. Oceanogr.*, 46, 1208–1214.
- Moustafa, A., Loram, J.E., Hackett, J.D., Anderson, D.M., Plumley, F.G. & Bhattacharya, D. (2009). Origin of saxitoxin biosynthetic genes in cyanobacteria. *PLoS ONE*, 4, e5758.
- Nielsen, L.T., Krock, B. & Hansen, P.J. (2013). Production and excretion of okadaic acid, pectenotoxin-2 and a novel dinophysistoxin from the DSP-causing marine dinoflagellate *Dinophysis acuta*: effects of light, food availability and growth phase. *Harmful Algae*, 23, 34–45.
- Orihel, D.M., Bird, D.F., Brylinsky, M., Chen, H.R., Donald, D.B., Huang, D.Y. *et al.* (2012). High microcystin concentrations occur only at low nitrogen-to-phosphorus ratios in nutrient-rich Canadian lakes. *Can. J. Fish. Aquat. Sci.*, 69, 1457–1462.
- Parkhill, J.P. & Cembella, A.D. (1999). Effects of salinity, light and inorganic nitrogen on growth and toxigenicity of the marine dinoflagellate *Alexandrium tamarense* from northeastern Canada. *J. Plankton Res.*, 21, 939–955.
- Quigg, A., Finkel, Z.V., Irwin, A.J., Rosenthal, Y., Ho, T.Y., Reinfelder, J.R. *et al.* (2003). The evolutionary inheritance of elemental stoichiometry in marine phytoplankton. *Nature*, 425, 291–294.
- Sanchez, S. & Demain, A.L. (2002). Metabolic regulation of fermentation processes. *Enzyme Microb. Technol.*, 31, 895–906.
- Scott, J.T., McCarthy, M.J., Otten, T.G., Steffen, M.M., Baker, B.C., Grantz, E.M. *et al.* (2013). Comment: an alternative interpretation of the relationship between TN:TP and microcystins in Canadian lakes. *Can. J. Fish. Aquat. Sci.*, 70, 1265–1268.
- Sivonen, K. & Jones, G. (1999). Cyanobacterial Toxins. In: *Toxic Cyanobacteria in Water: A Guide to their Public Health Consequences, Monitoring and Management* (eds Chorus, I., Bartram, J.). E & FN Spon, WHO London, pp. 41–112.
- Skovgaard, A., Legrand, C., Hansen, P.J. & Graneli, E. (2003). Effects of nutrient limitation on food uptake in the toxic haptophyte *Prymnesium parvum*. *Aquat. Microb. Ecol.*, 31, 259–265.
- Smalley, G.W., Coats, D.W. & Stoecker, D.K. (2003). Feeding in the mixotrophic dinoflagellate *Ceratium furca* is influenced by intracellular nutrient concentrations. *Mar. Ecol. Prog. Ser.*, 262, 137–151.
- Smith, V.H. (2003). Eutrophication of freshwater and coastal marine ecosystems: a global problem. *Environ. Sci. Pollut. Res.*, 10, 126–139.
- Staunton, J. & Weissman, K.J. (2001). Polyketide biosynthesis: a millennium review. *Natural Product Reports*, 18, 380–416.
- Sterner, R.W. & Elser, J.J. (2002). *Ecological Stoichiometry: The Biology of Elements from Molecules to the Biosphere*. Princeton University Press, Princeton.
- Stoecker, D.K., Tillmann, U. & Granéli, E. (2006). Phagotrophy in Harmful Algae. In *Ecology of Harmful Algae*. (eds Granéli, E., Turner, J.T.). Springer-Verlag, Berlin Heidelberg, Heidelberg, Germany, pp. 177–187.
- Sun, J., Hutchins, D.A., Feng, Y.Y., Seubert, E.L., Caron, D.A. & Fu, F.X. (2011). Effects of changing pCO<sub>2</sub> and phosphate availability on domoic acid production and physiology of the marine harmful bloom diatom *Pseudo-nitzschia multiseriata*. *Limnol. Oceanogr.*, 56, 829–840.
- Tillmann, U. (2003). Kill and eat your predator: a winning strategy of the planktonic flagellate *Prymnesium parvum*. *Aquat. Microb. Ecol.*, 32, 73–84.
- Tonk, L., Visser, P.M., Christiansen, G., Dittmann, E., Snelder, E., Wiedner, C. *et al.* (2005). The microcystin composition of the cyanobacterium *Planktothrix agardhii* changes toward a more toxic variant with increasing light intensity. *Appl. Environ. Microbiol.*, 71, 5177–5181.
- Touzet, N., Franco, J.M. & Raine, R. (2007). Characterization of nontoxic and toxin-producing strains of *Alexandrium minutum* (Dinophyceae) in Irish coastal waters. *Appl. Environ. Microbiol.*, 73, 3333–3342.
- Van de Waal, D.B., Verspagen, J.M.H., Lurling, M., Van Donk, E., Visser, P.M. & Huisman, J. (2009). The ecological stoichiometry of toxins produced by harmful cyanobacteria: an experimental test of the carbon-nutrient balance hypothesis. *Ecol. Lett.*, 12, 1326–1335.
- Wiedner, C., Visser, P.M., Fastner, J., Metcalf, J.S., Codd, G.A. & Mur, L.R. (2003). Effects of light on the microcystin content of *Microcystis* strain PCC 7806. *Appl. Environ. Microbiol.*, 69, 1475–1481.

## SUPPORTING INFORMATION

Additional Supporting Information may be downloaded via the online version of this article at Wiley Online Library ([www.ecologyletters.com](http://www.ecologyletters.com)).

Editor, James Grover

Manuscript received 21 October 2013

First decision made 27 November 2013

Second decision made 27 January 2014

Third decision made 13 March 2014

Manuscript accepted 17 March 2014